



## CUCUMBER FOLIAR FERTILIZATION (GREENHOUSE - FIELD)

PURPOSE OF USAGE	UTILIZATION TIME	PREMERGENCE INTO 100 LT WATER	BLOOMING PERIOD INTO 100 LT WATER	RIPENING PERIOD INTO 100 LT WATER	HARVEST PERIOD INTO 100 LT WATER
TO MEET THE NEED FOR PHOSPHORUS TO ENCOURAGE LATERAL GROWTH		VEJETOL 10-42-10 200 GR + FULL ENERGY 7 200 CC			
TO INHIBIT EXCESSIVE GROWTH TO DIMINISH THE DISTANCE BETWEEN NODES		FULLBASS 69 200-250 GR			
TO INCREASE NUMBER OF FLOWERS THE DEVELOP RESISTANCE TOWARDS PLANT DISEASES		FOLKAR-K 200-250 CC + NİMBUS 52 200-250 GR			
TO DIRECT THE PLANT TOWARDS FRUIT DEVELOPMENT TO HASTEN FRUIT DEVELOPMENT TO ELIMINATE Mn-Fe DEFICIENCIES TO CORRECT THE FRUIT AND LEAF COLOR			GLUKOMİX Min-Fe 200 CC + MAXİFORTE 250 GR		
TO PROMOTE FRUIT DEVELOPMENT TO ELIMINATE CALCIUM DEFICIENCY				MAXİFORTE 200-250 GR + CALVİN 200-250 CC	
TO ELIMINATE THE YELLOWING ARISING DUETO(N) TO PROMOTE VEGETATIVE GROWTH AND FRUIT				PROMAX COMBİ 80 GR + NİMBUS 52 200 GR	
TO PREVENT FRUIT DROWNING (FRUIT YELLOWING) TO ESTABLISH THE WATER BALANCE OF PLANT BY REGULATING PERSPIRATION					NOABSES 150 CC + CALVİN 200 CC