

SQUASH FOLIAR FERTTILIZATION (GREENHOUSE - FIELD)

UTILIZATION TIME PURPOSE OF USAGE	PREMERGENCEİ INTO 100 LT WATER	BLOMING AND RIPERNING PERIOD (DURING COLD PERIOD) INTO 100 LT WATER	RIPENING PERIOD INTO 100 LT WATER	RIPENING PERIOD INTO 100 LT WATER
TO PROMOTE THE PLANT DEVELOPMENT TO MEET THE Zn DEMAND TO PROMOTE LATERAL GROWTH AND CORE DEVELOPMENT	ZENİTH ₁ 150 CC NİMBUS 52 ₁ 200-250 GR FULL ENERGY 7 200 CC			
POLLINATION FLOWER POLLEN FORMATION TO ACCELERATE THE LONGITUDAL DEVELOPMENT OF FRUIT TO INCREASING THE NUMBER OF FLOWERS		GENESİS L26 + TRANS-POL 150 GR		
TO ACCELERATE THE DEVELOPMENT OF MANY FRUITS AT THE SAME TIME TO PROMOTE THE VEGETATIVE GROWTH TO ELIMINATE THE PLANT STRESES TO PROVIDE STANDART FRUIT FORMATION			NİMBUS 52 200-250GR + PROMAX COMBİ 100 GR	
TO CORRECT THE GROWTH OF SPIKES WHICH IS HARMED DUE TO EXCESS CHEMICAL USAGE TO ELIMINATE STRESS TO PROMOTE FLOWER AND FRUIT DEVELOPMENT				ZENİTH 150 CC + NOABSES 200 CC