

PEACH FOLIAR FERTILIZATION

UTILIZATION TIME PURPOSE OF USAGE	WHEN 15-20% BLOOMS INTO 100 LT WATER	WHEN 80% OF PETALS IS SHED INTO 100 LT WATER	WHEN EMBRIO STARTS HARDENING INTO 100 LT WATER
POLLINATION TO PROMOTE FRUIT GROWTH WITHOUT FECUNDATION TO ELIMINATE ABSCISSIC ACID ACCUMULATION ON THE FRUIT STEM DUE TO STRESS, AND, THUS, TO SUPPRESS FRUIT ABSCISSION	VEJETOL 29-5-10 200 GR + NOABSES 150 CC		
TO SUPPRESS FRUIT ABSCISSION TO ELIMINATE LEAF YELLOWING (TRACE ELEMENT DEFICIENCY) TO PROMOTE LEAF-OFFSHOOT DEVELOPMENT		NİMBUS 52 100 GR + FULL ENERGY 7 100 CC + GLUKOMİX Mn-Fe 200 CC	
TO GROW FRUIT HOMOGENOUSLY TO ENSURE ACCELERATED GROWTH OF FRUIT AND TO ENHANCE FRUIT QUALITY BY INCREASING WEIGHT (LARGE-SUCCULENT FRUIT)			MAXİFORTE 200 GR + NOABSES 150 CC
TO MEET THE NEED FOR POTASSIUM TO MAKE FRUIT LARGER TO HARVEST EARLY			VEJETOL 10-5-40 150 GR + POTANİA-NK 200-250 G