

## PICKLING CUCUMBER FOLIAR FERTILIZATION

PURPOSE OF USAGE	UTILIZATION TIME 6-7 DAYS AFTER FIELD PLANTING 1000 m <sup>2</sup>	WHEN FRUIT STARTS BEING VISIBLE 1000 m <sup>2</sup>	IN CASE OF HEAT STRESS 1000 m <sup>2</sup>	PERIOD WHEN VEGETATIVE GROWTH SLOWS DOWN 1000 m <sup>2</sup>
TO PROMOTE PLANT GROWTH TO ENHANCE BLOOMING TO ENSURE LATERAL GROWTH	VEJETOL 10-42-10 200 GR + FULL ENERGY 7 200 CC			
TO INCREASE NUMBER OF FRUITS AND TO ENHANCE FLOWERING		MAXIFORTE 200-250 GR		
TO PREVENT FRUIT LOSSES (FRUIT DROWNING AND YELLOWING) TO ENHANCE FRUIT DURABILITY TO MEET THE NEED FOR CALCIUM TO ADJUST WATER ORDER IN PLANT (TO ELIMINATE PHYSIOLOGICAL STRESSES)			NIMBUS 52 200 GR + CALVIN 200-250 CC	
TO PROMOTE VEGETATIVE GROWTH				NOABSES 200 CC + GLUKOMIX Mn-Fe 200 CC