



KIWI FOLIAR FERTILIZATION

UTILIZATION TIME PURPOSE OF USAGE	AT THE BEGINNING OF BUDBURST INTO 100 LT WATER	AT THE BEGINNING OF LEAF DEVELOPMENT INTO 100 LT WATER	AFTER FLOWERING UNTIL THE FRUIT IS CHICKPEA SIZE INTO 100 LT WATER	IMMEDIATELY AFTER THE FRUIT HALF SIZE INTO 100 LT WATER	PRE-HARVEST INTO 100 LT WATER	AFTER HARVEST INTO 100 LT WATER
FOR BACTERIAL AND FUNGAL DISEASES	SUPKOR CU 200 CC					
FOR FLOWER and SHOOT BUD FORMATION FOR HEALTHY FRUIT SET	2000 ZENITH 150 cc + TRANSPOL 150 cc	FULLBASS 69 250 GR + TRANSPOL 150 GR + 2000 AGRİ BOLBOR 150 cc				
TO REDUCE FRUIT DROP			VEJETOL 29-5-10 200 GR + GENESIS L26 200 CC + CALVIN 250 CC			
TO ENSURE LARGER SIZED FRUIT TO IMPROVE THE QUALITY AND WEIGHT OF FRUIT TO ENSURE HOMOGENOUS ENLARGEMENT				MAXIFORTE 200-250 GR + NOABSES 100 CC + CALVIN 250 CC		
FOR FRUIT COLOR, FLAVOR, TASTE AND SHELF LIFE					ULTRA K30 300 CC + VEJETOL 10-5-40 200 GR	
FOR THE FORMATION OF HEALTHY FLOWER AND SHOOT BUD FOR THE FOLLOWING YEAR FOR BACTERIAL AND FUNGAL DISEASES						FOLKAR-K 250 CC + PROLIUM 250 CC