

HAZELNUT FOLIAR FERTILIZATION UTILIZATION TIME

UTILIZATION TIME PURPOSE OF USAGE	AT THE BEGINNING OF BUDBURST	AT THE BEGINNING OF LEAF DEVELOPMENT	AFTER FLOWERING UNTIL THE FRUIT IS DEVELOPMENT INTO 100 LT WATER	FRUIT ENLARGEMENT STAGE INTO 100 LT WATER	PRE-HARVEST INTO 100 LT WATER	AFTER HARVEST INTO 100 LT WATER
	INTO 100 LT WATER	INTO 100 LT WATER	INTO TOO EI WATER	INTO 100 EI WATER		
FOR BACTERIAL AND FUNGAL DISEASES	SUPKOR CU 200 CC					
FOR HEALTHY BUD AND FLOWER FORMATION	2000 ZENİTH 150 cc + TRANSPOL 150 cc					
FOR HEALTHY POLLINATION AND FRUIT SET		FULBAS 69 69 250 GR TRANSPOL 150 GR 2000 AGRİ BOLBOR 150 cc	GENESİS L26 200 CC CALVİN ⁺ 250 CC			
TO REDUCE FRUIT DROP AND TO Ensure homogeneous set of fruits				MAXIFORTE 200-250 GR NOABSES 100 CC CALVIN ⁺ 250 CC		
TO ENCOURAGE LARGER SIZED FRUITS WITH SUCCULENT CORES					ULTRA K30 300 CC VEJETOL 10 ⁻⁵ -40 200 GR	
FOR THE FORMATION OF HEALTHY FLOWER And shoot bud for the following year for bacterial and fungal diseases						SUPKOR CU 200 CC 7-10 GÜN SONRA 2000 ZENİTH 150 CC + TRANSPOL 150 GR