



WALNUT FOLIAR FERTILIZATION UTILIZATION TIME

PURPOSE OF USAGE \ UTILIZATION TIME	AT THE BEGINNING OF BUDBURST INTO 100 LT WATER	AT THE BEGINNING OF LEAF DEVELOPMENT INTO 100 LT WATER	AFTER FLOWERING UNTIL THE FRUIT IS CHICKPEA SIZE INTO 100 LT WATER	IMMEDIATELY AFTER THE FRUIT HALF SIZE INTO 100 LT WATER	AFTER HARVEST INTO 100 LT WATER
FOR BACTERIAL AND FUNGAL DISEASES	SUPKOR CU 200 CC				
FOR HEALTHY BUD AND FLOWER FORMATION	2000 ZENITH 150 cc + TRANSPOL 150 cc	FULBAS 69 69 250 GR + TRANSPOL 150 GR + 2000 AGRİ BOLBOR 150 cc			
FOR HEALTHY POLLINATION AND FRUIT SET			MAXIFORTE 200-250 GR + NOABSES 100 CC + CALVİN 250 CC		
TO REDUCE FRUIT DROP AND TO ENSURE HOMOGENEOUS SET OF FRUITS				ULTRA K 30 300 CC + VEJETOL 10-5-40 200 GR	
TO ENCOURAGE LARGER SIZED FRUITS WITH SUCCULENT CORE					FOLKAR-K 250 CC + PROLIUM 250 CC