

STRAWBERRY FOLIAR FERTILIZATION (GREENHOUSE)

5	UTILIZATION TIME PURPOSE OF USAGE	10-15 DAYS POST EMERGENCE INTO 100 LT WATER	WHEN PEDICLE STARTS GROWING INTO 100 LT WATER	PREMERGENCE INTO 100 LT WATER	WHEN FRUIT STARTS DEVELOPING INTO 100 LT WATER	IF VEGETATIVE GROWTH IS SLOW INTO 100 LT WATER
	TO INCREASE THE VEGETATIVE GROWTH TO MEET THE Mn-Fe DEMAND	GLUKOMİX Mn-Fe 200 CC + FULL ENERGY 7 200 CC		- A		
	TO PROMOTE DEVELOPMENT TO ELIMINATE THE NEED FOR TRACE ELEMENT	PROMAX COMBİ 80 GR + NİMBUS 52 200 GR				
	TO OBTAIN ACCELARATED DEVELOPMENT OF PEDICLES TO INCREASE THE NUMBER OF PEDICLES (YIELD ENHANCEMENT)		PROMAX COMBİ 100 GR + GENESİS L26 200-250 C			
	TO PROMOTE FLOWERING POLLINATION			ZENİTH 150 CC , GENESİS L26 200 CC		
	TO PROMOTE FRUIT DEVELOPMENT TO ENHANCE COLOR AND QUALITY TO ELIMINATE CALCIUM DEFICIENCY				MAXIFORTE 250 GR + CALVIN 250 CC	
	TO PROMOTE VEGETATIVE GROWTH TO MEET THE NEED FOR TRACE ELEMENT					GLUKOMİX Mn-Fe 200 cc + NİMBUS 52 200 gr