

## TOMATO FOLIAR FERTILIZATION (GREENHOUSE-FIELD)

PURPOSE OF USAGE	UTILIZATION TIME	SEEDLING PERIOD INTO 100 LT WATER	PREMERGENCE WHILE CLUSTER IS BEING FORMED INTO 100 LT WATER	WHEN FRUIT START TO EMERGE INTO 100 LT WATE	WHEN FRUIT IS HALF-SIZE INTO 100 LT WATER	AT THE BEGINNING OF COLORING INTO 100 LT WATER
TO INHIBIT EXCESSIVER GOWTH (TO DIMINISH THE DISTANCE BETWEEN NODES) TO PROMOTE THE PLANT TOWARDS ROOT DEVELOPMENT TO DEVELOP RESISTANCE TOWARD DISEASES		FOLKAR-K 200-250 CC + MAXIFORTE 200 GR				
TO DIMINISH THE DISTANCE BETWEEN NODES			FULLBASS 69 200-250 GR			
TO ENHANCE FLOWERING AND POLLEN FORMATION BY ELEMINATING STRESSES TO PROVENT FLOWER ABCISSION			ZENİTH 150 CC + NOABSES 200 CC			
TO ACCELERATE FRUIT AND FLOWER DEVELOPMENT TO REGULATE NUTRIENT UP-TAKE THROUGH ROOT SYSTEM AND KEEP THE HORMON ANZIME BALANCE IN ORDER				ZENİTH 150 CC + MAXIFORTE 200 GR		
TO OBTAIN LARGER FRUITS AND IN COLD SEASONS,IT PREVENT CHLOROSIS					NOABSES 200 CC + CALVİN 200-300 CC	
TO OBTAIN LARGER FRUITS AND IN COLD SEASONS,IT PREVENT CHLOROSIS					MAXIFORTE 200 GR + GLUKOMİX Mn-Fe 200 cc	
WHEN,FRUIT DEVELOPMENT AND FLOWERING OCCURE AT THE SAME TIME, TO PREVENT B DEFICIENCY TO SUPPORT POLLEN FORMATION,AND TO ACCELERATE VEGETATIVE GROWTH					TRANS-POL 150 GR + NOABSES 200 CC	
IT PROVIDES FAST GROWTH OF FRUITS AND HASTEN HOMOGENOUS COLORING						MAXIFORTE 200 GR + VEJETOL 10-5-40 200 GR + GLUKOMİX Mn-Fe 200 CC